

**(REVISED COURSE)**

Q P Code : NP-17690

(3 Hours)

[Total Marks : 80]

- N.B. : (1) Questions No. 1 is compulsory.  
 (2) Attempt any three from the remaining questions.  
 (3) Assume suitable data if necessary.

1. (a) Prove that  $\operatorname{Sech}^{-1}(\sin\theta) = \log\left(\cot\frac{\theta}{2}\right)$  3

(b) If  $x = \cos\theta - r\sin\theta$ ,  $y = \sin\theta + r\cos\theta$  3  
 prove that  $\frac{dr}{dx} = \frac{x}{r}$

(c) If  $x = e^v \sec u$ ,  $y = e^v \tan u$  3  
 find  $J\left(\frac{u, v}{x, y}\right)$

(d) If  $y = \sin px + \cos px$  3  
 Prove that  $y_n = p^n [1 + (-1)^n \sin(2px)]^{\frac{1}{2}}$

(e) Find the series expansion of  $\log(1+x)$  in powers of  $x$ . Hence prove that 4  
 $\log x = (x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{3}(x-1)^3 \dots$

(f) If 'A' is skew-symmetric matrix of odd order then prove that it is singular. 4

2. (a) Show that the roots of the equation  $(x+1)^6 + (x-1)^6 = 0$  are given by 6

$$-i\cot\left(\frac{2n+1}{12}\pi, n=0,1,2,3,4,5\right)$$

(b) Find two non-singular matrices P & Q such that PAQ is in normal form where 6

$$A = \begin{bmatrix} 1 & 2 & 3 & -4 \\ 2 & 1 & 4 & -5 \\ -1 & -5 & -5 & 7 \end{bmatrix}$$

Also find rank of A.

(c) If  $x + y = 2e^\theta \cos\phi$ ,  $x - y = 2ie^\theta \sin\phi$  & u is a function of x & y then prove that 8

$$\frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial \phi^2} = 4xy \frac{\partial^2 u}{\partial x \partial y}$$

3. (a) Find the value of  $\lambda$  for which the equations  $x_1 + 2x_2 + x_3 = 3$ ,  $x_1 + x_2 + x_3 = \lambda$ ,  $3x_1 + x_2 + 3x_3 = \lambda^2$  has a solution & solve them completely for each value of  $\lambda$ . 6
- (b) Divide 24 into three parts such that the product of the first, square of the second & cube of the third is maximum. 6
- (c) (i) If  $\operatorname{cosec}\left(\frac{\pi}{4} + ix\right) = u + iv$  prove that  $(u^2 + v^2)^2 = 2(u^2 - v^2)$  4  
(ii) Prove that  $\tan\left(i \log\left(\frac{a - ib}{a + ib}\right)\right) = \frac{2ab}{a^2 - b^2}$  4
4. (a) Show that  $\frac{\partial(u, v)}{\partial(x, y)} = 6r^3 \sin 2\theta$  given that  $u = x^2 - y^2$ ,  $v = 2x^2 - y^2$  &  
 $x = r \cos \theta$ ,  $y = r \sin \theta$ . 6
- (b) If  $\alpha = 1 + i$ ,  $\beta = 1 - i$  &  $\cot \theta = x + 1$   
prove that  $(x + \alpha)^n + (x + \beta)^n = (\alpha + \beta) \cos n\theta \operatorname{cosec}^n \theta$ . 6
- (c) Using Gauss-seidel method, solve the following system of equations upto 3<sup>rd</sup> iteration. 8  

$$\begin{aligned} 5x - y &= 9 \\ -x + 5y - z &= 4 \\ -y + 5z &= -6 \end{aligned}$$
5. (a) Using De-Moivr's theorem, prove that 6  

$$\frac{\sin 6\theta}{\sin \theta} = 16 \cos^4 \theta - 16 \cos^2 \theta + 3$$
- (b) Explain  $\frac{x}{e^x - 1}$  in powers of  $x$ . 6  
Hence prove that  $\frac{x}{2} \left[ \frac{e^x + 1}{e^x - 1} \right] = 1 + \frac{1}{12}x^2 - \frac{1}{720}x^4 + \dots$
- (c) If  $y = \frac{\sin^{-1} x}{\sqrt{1 - x^2}}$  8  
prove that  $(1 - x^2)y_{n+2} - (2n + 3)xy_{n+1} - (n + 1)^2 y_n = 0$ . Hence find  $y_n(0)$

**Con. 11513-14.****TURN OVER**

6. (a) Examine the linear dependence or independence of vectors  $(1, 2, -1, 0)$ ,  $(1, 3, 1, 3)$ ,  $(4, 2, 1, -1)$  &  $(6, 1, 0, -5)$  6

(b) If  $u = f\left(\frac{x-y}{xy}, \frac{z-x}{xz}\right)$  prove that 6

$$x^2 \frac{\partial u}{\partial x} + y^2 \frac{\partial u}{\partial y} + z^2 \frac{\partial u}{\partial z} = 0$$

(c) (i) Fit a straight line to the following data with  $x$  - as independent variable. 4

X : 1965 1966 1967 1968 1969

Y : 125 140 165 195 200

(ii) Evaluate  $\lim_{x \rightarrow 0} (1 + \tan x)^{\cot x}$  4