

ExTC/Inst/A-M.IV/IV/61/1216

App. Maths - IV / 01/12/16

Q. P. Code : 545800

Duration: 3 Hours

(Revised Course)

Total Marks: 80



N.B. : 1) Q.1. is compulsory.

2) Attempt any three from the remaining.

Q.1. a) If $f(x)$ is an algebraic polynomial in x and λ is an eigen value and X is the corresponding eigen vector of a square matrix A then $f(\lambda)$ is an eigen value and X is the corresponding eigenvector of $f(A)$. (5)

b) Find the extremal of $\int_{x_0}^{x_1} (x + y')y' dx$ (5)

c) Express (6,11,6) as linear combination of $v_1 = (2,1,4), v_2 = (1,-1,3), v_3 = (3,2,5)$. (5)

d) Evaluate $\int_C \frac{z}{(z-1)^2(z-2)} dz$, where C is the circle $|z-2|=0.5$ (5)

Q.2. a) Find the curve $y = f(x)$ for which $\int_0^{\pi} (y'^2 - y^2) dx$ is extremum if $\int_0^{\pi} y dx = 1$. (6)

b) Evaluate $\int_0^{2\pi} \frac{\cos 3\theta}{5 + 4 \cos \theta} d\theta$ (6)

c) Find the singular value decomposition of $\begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$ (8)

Q.3. a) Verify Cayley Hamilton theorem for $A = \begin{bmatrix} 3 & 10 & 5 \\ -2 & -3 & -4 \\ 3 & 5 & 7 \end{bmatrix}$ and hence, find the matrix represented by $A^6 - 6A^5 + 9A^4 + 4A^3 - 12A^2 + 2A - I$. (6)

b) Construct an orthonormal basis of R^3 using Gram Schmidt process to $S = \{(3,0,4), (-1,0,7), (2,9,11)\}$ (6)

c) Find all possible Laurent's expansions of $\frac{z}{(z-1)(z-2)}$ about $z = -2$ indicating the region of convergence. (8)

[Turnover]

Q.4. a) Reduce the quadratic form $2x^2 - 2y^2 + 2z^2 - 2xy - 8yz + 6zx$ to canonical form and hence, find its rank, index and signature and value class. (6)

b) If $\phi(\alpha) = \int_C \frac{4z^2 + z + 5}{z - \alpha} dz$, where C is the contour of the ellipse $\frac{x^2}{4} + \frac{y^2}{9} = 1$, find the values of $\phi(3.5), \phi(i), \phi'(-1), \phi''(-i)$ (6)

c) Using Rayleigh-Ritz method, solve the boundary value problem $I = \int_0^1 (y'^2 - y^2 - 2xy) dx$; $0 \leq x \leq 1$, given $y(0) = y(1) = 0$. (8)

Q.5. a) Find the extremal of the function $\int_0^{\pi/2} (2xy + y^2 - y'^2) dx$; with $y(0) = 0, y(\pi/2) = 0$ (6)

b) Find the orthogonal matrix P that diagonalises $A = \begin{bmatrix} 4 & 2 & 2 \\ 2 & 4 & 2 \\ 2 & 2 & 4 \end{bmatrix}$ (6)

c) Using Cauchy's Residue theorem, evaluate $\oint_C \frac{z^2 + 3}{z^2 - 1} dz$ where C is the circle (i) $|z - 1| = 1$ (ii) $|z + 1| = 1$. (8)

Q.6. a) Find the sum of the residues at singular points of $f(z) = \frac{z}{(z-1)^2(z^2-1)}$ (6)

b) If $A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$, prove that $A^{50} - 5A^{49} = \begin{bmatrix} 4 & -4 \\ -2 & 2 \end{bmatrix}$ (6)

c) (i) Check whether $W = \{(x, y, z) | y = x + z, x, y, z \text{ are in } \mathbb{R}\}$ is a subspace of \mathbb{R}^3 with usual addition and usual multiplication. (4)

(ii) Find the unit vector in \mathbb{R}^3 orthogonal to both $u = (1, 0, 1)$ and $v = (0, 1, 1)$. (4)